

Chapter 10

Hand Hygiene

Carolina Giuffré and Claire Kilpatrick

Key Points

- Hand hygiene is the most effective single measure to prevent transmission of healthcare-associated pathogens.
- Compliance with hand hygiene recommendations is often sub-optimal; compliance is influenced by many factors, including equipment/supplies, time constraints, and behavioural factors.
- Hand hygiene can be performed either rubbing with an alcohol-based hand rub or by washing with soap and water. The World Health Organization recommends the preferred use of alcohol-based hand rub for routine hand hygiene in health care, if available.
- Hand hygiene promotion and multimodal improvement strategies have a great impact on healthcare worker practices and can reduce healthcare-associated infections and the spread of resistant microorganisms.
- Effective strategies include: provision of alcohol-based hand rubs and clean water, soap, and disposable towels; staff education; monitoring of hand hygiene practices and performance feedback; reminders in the work place; and promotion of a patient safety climate.

Background

Hand washing with soap and water has been considered a measure of personal hygiene for centuries. In the mid-1800s, studies by Ignaz P. Semmelweis in Vienna, Austria, and Oliver Wendell Holmes in Boston, Massachusetts, USA, established that hospital-acquired infections were transmitted via the hands of healthcare workers (HCW).¹ Following the observation of high maternal mortality rates due to puerperal fever, Semmelweis made physicians wash their hands in a chlorinated lime solution before every patient contact. Knowledge of the transmission of pathogens through hands and of infection prevention has greatly evolved, and the first international guidelines on hand hygiene published in 2009 recommended a range of evidence-based actions.¹

Resident or transient microbial flora is present on normal human skin. Resident flora are protective and less likely to be associated with healthcare-associated infections (HAI), however they can cause contamination of sterile body cavities, eyes, or non-intact skin. Transient flora colonise the superficial layers of the skin and may cause HAIs as they are acquired and passed on by HCWs during direct contact with patients or contaminated environmental surfaces. Contaminated HCWs' hands are the commonest route of transmission of HAIs. Hand hygiene is therefore the most effective measure to prevent HAIs.¹

Hand Contamination

HCWs hands can become contaminated, even if gloved, with pathogens, such as *Staphylococcus aureus*, enterococci, *Clostridium difficile*, Gram-negative bacilli, and some viruses (e.g., respiratory syncytial virus and rhinovirus). This can happen by touching patient skin or body sites contaminated or infected with pathogens (known or unknown) including draining wounds, as well as contaminated surfaces within the health care environment, especially surrounding the patient.¹ Some activities in particular can lead to heavier hand contamination and the potential for pathogen spread if the right actions are not taken. Examples include direct patient touch, touching body fluids or waste, diaper change, and respiratory care. In addition, HCWs with dermatitis or skin lesions on their hands may remain colonised with acquired microorganisms for a significant period of time.¹

Subungual (beneath the nails) areas of the hand carry high concentrations of microorganisms, contributing to the spread of pathogens. Artificial nails may also contribute to the transmission of pathogens as wearers are more likely to harbour Gram-negative bacilli on their fingertips than those with natural nails, despite hand washing or rubbing with an alcohol-based product. Diseased fingernails reduce the efficacy of hand hygiene.¹ The skin underneath rings (including wedding rings) is more heavily colonised than that on other fingers. Rings with sharp and voluminous surfaces and long, sharp fingernails, either natural or artificial, can puncture gloves and limit HCWs' hand hygiene efficacy.¹

Compliance Among Healthcare Workers

Without adequate hand hygiene, hand contamination increases; contaminated HCWs' hands have been associated with endemic HAIs.¹ Therefore, hand hygiene is the primary measure to prevent HAIs and will also help decrease the spread of multidrug resistant microorganisms, thus supporting the current global antimicrobial resistance agenda. However, many determinants, such as lack of time, lack of equipment/supplies, and behavioural factors, often result in HCWs neglecting hand hygiene. Although many HCWs perceive their performance as high,²⁻⁵ their hand hygiene compliance is usually < 40% in the absence of interventions.^{1,6}

Hand hygiene performance varies according to work intensity, type of ward, professional category, and time of day/week. Compliance can be lower in settings with high care intensity (e.g., intensive care units), among physicians, and before rather than after touching a patient.^{1,6} Indeed, HCWs tend to comply more frequently with indications that protect themselves (e.g., after exposure to body fluids, after glove use, after

contact with the patient or the patient's environment).^{1,7} This is important in understanding how to affect behaviours and ensure patient safety in relation to HAI prevention.

Products and Techniques

Hand hygiene can be performed either by rubbing with an alcohol-based formulation or by washing with soap and water. Liquid soap is preferred; if there is no other choice, bar, leaf, or powder may be used. Supplies must be placed alongside running water and re-stocked when needed in order to achieve compliance. Plain soap has minimal antimicrobial activity, nonetheless it is recommended for hand washing because mechanical friction alone removes many transient microorganisms. (See Table 10.1)^{1,8}

The commonest antimicrobials in hand hygiene products are: alcohols, chlorhexidine, iodine, and iodophors. Studies suggest efficacy against Gram-positive and Gram-negative bacteria with maximum efficacy is demonstrated by alcohols and iodophors.¹ Mycobacteria and fungi are most effectively eliminated by alcohols, less so by chlorhexidine.

Enveloped viruses (e.g., herpes simplex, human immunodeficiency, influenza, and respiratory syncytial) are highly susceptible to alcohols;^{1,9} hepatitis B and C viruses require high concentrations (70-80% [volume/volume (v/v)]).¹ Alcohols have shown *in vivo* activity against some non-enveloped viruses (rotavirus, adenovirus, rhinovirus, hepatitis A virus, and enteroviruses).¹ *In vitro* virucidal activity against surrogate strains of norovirus was demonstrated by 70% alcohol-based formulations. Several norovirus outbreaks were controlled with preventive measures, including use of alcohol-based hand rub.^{1,10-11} In general, ethanol has greater activity against viruses than isopropanol.

Iodophors and chlorhexidine have some activity against enveloped and some non-enveloped viruses.¹ None of these antiseptics has activity against bacterial spores or protozoan oocysts, although the mechanical effect of washing with soap and water allows their partial removal.^{1,8,12}

According to the World Health Organization (WHO) alcohol-based hand rubs should be the preferred method for hand hygiene. See Table 10.1.^{1,8} These products have the broadest antimicrobial spectrum, require a short time (20-30 sec) for effective antimicrobial decontamination, have better skin tolerance, and are readily available at the point of care (i.e., where care is provided).^{1,13} See Figure 10.1.




Figure 10.1. Time-course of efficacy of unmedicated soap and water and alcohol-based hand rub in reducing the release of test bacteria from artificially-contaminated hands. (Reprinted from *Lancet Inf Dis* 2001; Suppl 1:14 with permission from Elsevier)

The efficacy of an alcohol-based hand rub depends on its quality, the amount used, the time spent rubbing, and complete coverage of the hands' surfaces (See Figure 10.2). These parameters also apply to washing with soap and water (See Figure 10.3). Hand rubs containing 60–80% alcohol are satisfactory, provided that they meet recommended standards (European Norms [EN] or American Society for Testing and Materials [ASTM] standards).^{1,14} 75-87% ethanol, isopropanol, or n-propanol, or a combination of these products guarantee the optimal antimicrobial efficacy. The WHO-recommended formulations contain either 75% v/v isopropanol or 80% v/v ethanol.^{1,15} There is clear evidence that local manufacture of quality products is possible in low resource settings.¹⁶ This formulation is listed in the WHO Essential Medicines List.

It is worth noting that during the Ebola virus disease outbreak of 2014-2015 in West Africa, the use of chlorine for hand washing was prominently featured. Despite some statements that it was part of outbreak control, there is emerging evidence of the harm caused by its over and misuse. WHO published a guideline recommending that due to lack of evidence, health facilities should move to the use of alcohol hand rub and soap and water over the use of chlorine.¹⁷



Figure 10.2. Hand hygiene technique with an alcohol-based formulation (Based on the hand hygiene technique with an alcohol-based formulation, http://www.who.int/gpsc/5may/tools/system_change/en/index.html © World Health Organization 2009. All rights reserved.)

Alcohol-based hand rubs are available as rinses (with low viscosity), gels, foams, and impregnated wipes. However, the wipes require more evidence before they can be used effectively in practice.¹⁸⁻²⁰

Points to consider when selecting a product include:^{1,15}

- 1) demonstrated antimicrobial efficacy according to ASTM or EN standards for hygienic hand antisepsis and/or surgical hand preparation;
- 2) proven good dermal tolerance and minimal skin reactions;
- 3) minimum drying time (products that require longer drying times may affect hand hygiene best practice);
- 4) cost;
- 5) aesthetic preferences of HCWs and patients, such as fragrance, colour, texture, “stickiness”, and ease of use; and
- 6) availability, convenience, and functioning of dispensers, and ability to prevent their contamination.¹³

A rational location of facilities and supplies (i.e., sinks, soap, towels, and hand rub dispensers), as well as good maintenance and user-friendliness, are essential to facilitate hand hygiene when it is most needed at the point of care. Ideally, different alcohol-based hand rub dispensers, e.g., pocket bottles, wall-mounted, or those placed on carts/trolleys, night stand/bedside table, or affixed to the bed rail, should be used. Confirmation of the need to extend the use of hand rub at the point where care is provided was published by the WHO in its report on water, sanitation and hygiene in health care facilities.²¹⁻²² It states that 35% of facilities do not have water and soap for handwashing.

There is a misconception that handwashing must be followed by use of an alcohol based hand rub for anti-sepsis. The recommendation not to do this is clearly featured in the WHO Guidelines¹ and is an important resource efficient consideration.



Figure 10.3. Hand washing technique with soap and water (Based on the hand washing technique with soap and water, [URL:http://www.who.int/gpsc/5may/tools/system_change/en/index.html](http://www.who.int/gpsc/5may/tools/system_change/en/index.html) © World Health Organization 2009. All rights reserved.)

When to Perform Hand Hygiene

The “My five moments for hand hygiene” approach^{1, 23-24} (See Figure 10.4) merges the hand hygiene indications¹ into five moments when hand hygiene is required, focused on microbial risk as well as human factors thinking. These are:

- 1) Before touching a patient,
- 2) Before clean/aseptic procedures,
- 3) After body fluid exposure/risk,
- 4) After touching a patient, and

5) After touching patient surroundings.

This approach proposes a unified vision for HCWs, trainers, and observers to minimise inter-individual variation in practice.



Figure 10.4. “My five moments for hand hygiene” (reproduced with permission from reference 12)

Glove Use

Gloves are used as personal protective equipment (PPE) to protect HCWs’ hands. They can reduce transmission of pathogens and help control outbreaks as part of PPE recommended use. However, gloves are only effective in prevention of HAI if rigorously accompanied by other measures, including hand hygiene; the use of gloves never negates the need for hand hygiene.¹ To prevent HCW contamination, gloves must also be used according to established indications for donning and removal.

Use of the same gloves for several hours, while providing care to different patients and touching multiple surfaces, is a very frequent malpractice. Several studies have established an association between inappropriate glove use and low compliance with hand hygiene. Others have found that HCWs wearing gloves were significantly more likely to cleanse their hands following patient care.^{25,26}

Understanding that glove use does not replace hand hygiene is of utmost importance. When there is a need for performing hand hygiene (opportunity) before a care act which also requires glove use (indicating Moment 2 of the WHO 5 Moments), hand rubbing or handwashing must be performed before donning gloves, as well as immediately after glove removal.²⁷ Within the ‘patient zone’ if a Moment 2 or 3 arises, whether gloves are worn or not, hand hygiene should be performed, with removal of gloves at these times.²⁸⁻³⁰

Improvement Strategies¹

Key components of successful strategies are:

System change

Ensure that the necessary infrastructure is in place to allow HCWs to practice hand hygiene. This includes two essential elements:

- 1) provision of alcohol-based hand rub at the point of care; and

- 2) access to a safe, continuous, water supply, soap, and disposable towels.

Training/education

Provide regular training on microbial transmission through HCWs' hands and the importance of hand hygiene based on the "My five moments for hand hygiene" approach. Also include the correct procedures for hand rubbing and hand washing by using real life clinical scenarios in presentations, e-learning modules, posters, focus groups, reflective discussion, videos, self-learning modules, practical demonstrations, feedback from assessment, buddy systems, or combinations of these methods. Assess the impact of training on HCWs' knowledge to identify areas for further education.

Evaluation and feedback

Monitor hand hygiene practices and knowledge among HCWs and provide results to staff. The gold standard for measuring hand hygiene compliance remains direct observation; electronic monitoring of hand hygiene actions and evaluation of alcohol-based hand rub consumption can be used as indirect methods and surrogate markers.³¹ The applicability of electronic systems needs serious consideration given the associated costs and the actual potential for behaviour change. Use of the WHO Hand Hygiene Self-Assessment Framework provides a score to use for evaluating facility progress with hand hygiene improvement.²⁹⁻³⁰

Reminders in the workplace

Remind HCWs about the importance of hand hygiene and the indications and procedures for performing it. Change these prompts on a regular basis to continue to have an impact on behaviour.

Institutional safety climate

Create an environment that raises awareness about patient safety while making hand hygiene a high priority at all levels, including active participation at institutional and individual levels.

Elements of this multimodal strategy to improve hand hygiene demonstrated reduction in HAI and cross-transmission rates of potential pathogens.^{1,32-33} Multimodal interventions are considered most effective for compliance.³⁴⁻³⁵

It has become increasingly clear that hand hygiene is a behaviourally driven science. Hand hygiene is a *modifiable* behavioural risk factor for HAI. Goal setting, award incentives, and accountability have recently been highlighted as important elements of achieving a safety climate. Addressing different combinations of interventions is deemed important.^{32,35-37} Human factors could play a role in behaviour change and compliance if the IPC community engages with other disciplines such as social sciences.³⁸⁻⁴¹

Summary

HCWs' hands play a crucial role in the transmission of microorganisms during the sequence of care and contact with environmental surfaces and patients' skin. Hand hygiene is the single most effective measure to prevent HAI. However, hand hygiene practice at the right moment using proper technique is usually sub-optimal among HCWs due to many perceived constraints and behavioural factors. Improvement of practices can be achieved and lead to substantial reduction of HAI by multimodal behaviour change strategies aimed at strengthening infrastructure, knowledge, and the institutional patient safety culture. Hand hygiene is noted as one of the Top Patient Safety Strategies that can be encouraged for adoption.⁴² The preferred use of alcohol-based hand rubbing as the gold standard for hand hygiene and the identification of the right moments for hand hygiene during patient care are essential elements for success.

Acknowledgement

This chapter is an update of the earlier one by Dr Benedetta Allegranzi, Professor Didier Pittet, and Claire Kilpatrick.

References

1. World Health Organization. WHO Guidelines on Hand Hygiene in Health Care. Geneva: World Health Organization; 2009. http://whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf [Accessed 22 February 2016].
2. Allegranzi B, Sax H, Bengaly L, et al. Successful implementation of the World Health Organization hand hygiene improvement strategy in a referral hospital in Mali, Africa. *Infect Control Hosp Epidemiol* 2010; 31:133-41.
3. Burnett E. Perceptions, attitudes, and behavior towards patient hand hygiene. *Am J Infect Control* 2009; 37:638-42.
4. Tai JW, Mok ES, Ching PT, Seto WH, Pittet D. Nurses and physicians' perceptions of the importance and impact of healthcare-associated infections and hand hygiene: a multi-center exploratory study in Hong Kong. *Infection* 2009; 37:320-33.
5. Hosseinalhashemi M, Sadeghipour Kermani F, Palenik CJ, Pourasghari H, Askarian M. Knowledge, attitudes, and practices of health care personnel concerning hand hygiene in Shiraz University of Medical Sciences hospitals, 2013-2014. *Am J Infect Control* 2015; 43 (9): 1009-1011.
6. Erasmus V, Daha TJ, Brug H, et al. Systematic review of studies on compliance with hand hygiene guidelines in hospital care. *Infect Control Hosp Epidemiol* 2010; 31:283-94.]
7. Borg MA, Benbachir M, Cookson BD, et al. Self-protection as a driver for hand hygiene among healthcare workers. *Infect Control Hosp Epidemiol* 2009; 30:578-80.
8. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings. Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *Morb Mortal Wkly Rep Recomm* 2002 25; 51:1-45. <http://www.cdc.gov/mmwr/PDF/rr/rr5116.pdf> [Accessed 22 February 2016]
9. Grayson ML, Melvani S, Druce J, et al. Efficacy of soap and water and alcohol-based hand-rub preparations against live H1N1 influenza virus on the hands of human volunteers. *Clin Infect Dis* 2009; 48:285-91.
10. Cheng VC, Tai JW, Ho YY, Chan JF. Successful control of norovirus outbreak in an infirmary with the use of alcohol-based hand rub. *J Hosp Infect* 2009; 72:370-1.
11. Centers for Disease Control and Prevention. Norovirus outbreaks on three college campuses - California, Michigan, and Wisconsin, 2008. *Morb Mortal Wkly Rep* 2009; 58:1095-100.
12. Oughton MT, Loo VG, Dendukuri N, Fenn S, Libman MD. Hand hygiene with soap and water is superior to alcohol rub and antiseptic wipes for removal of *Clostridium difficile*. *Infect Control Hosp Epidemiol* 2009; 30:939-44.
13. World Health Organization. Alcohol-based handrub planning and costing tool 2010. http://www.who.int/gpsc/5may/tools/system_change/en/index.html [Accessed 22 February 2016]
14. Rotter M, Sattar S, Dharan S, Allegranzi B, Mathai E, Pittet D. Methods to evaluate the microbicidal activities of hand-rub and hand-wash agents. *J Hosp Infect* 2009; 73:191-9.
15. World Health Organization. Guide to local production: WHO-recommended handrub formulations. 2010. http://www.who.int/gpsc/5may/Guide_to_Local_Production.pdf [Accessed 22 February 2016]
16. Bauer-Savage J., Pittet D., Kim E., Allegranzi B. Local production of WHO-recommended alcohol-based handrubs: feasibility, advantages, barriers and cost. <http://www.who.int/bulletin/volumes/91/12/12-117085/en/> [Accessed 22 February 2016]
17. WHO Guideline on hand hygiene in Health Care in the context of Filovirus disease outbreak response.

- Rapid advice guideline. November 2014. <http://www.who.int/csr/resources/publications/ebola/hand-hygiene/en/> [Accessed 22 February 2016]
18. Kampf G, Marschall S, Eggerstedt S, Ostermeyer C. Efficacy of ethanol-based hand foams using clinically relevant amounts: a cross-over controlled study among healthy volunteers. *BMC Infect Dis* 2010; 10:78.
 19. Butz A. et col. Alcohol-Impregnated wipes as an alternative in hand hygiene. *Amer J Infect Control* 1990; 18: 70-76.
 20. D'Antonio N. et col. Revisiting the hand wipe versus gel rub debate: Is a higher-ethano content hand wipe more effective than an etanol gel rub? *Amer J Infect Control* 2010; 38: 678-682.
 21. WHO (2015) Partnerships for Safer Healthcare Delivery: Evaluation of WHO African Partnerships for Patient Safety 2009-2014. <http://www.who.int/patientsafety/implementation/apps/evaluation-report/en/> [Accessed 22 February 2016]
 22. WHO Water, Sanitation and Hygiene in Health Care Facilities Status on low and middle-income countries and way forward, 2015. http://www.who.int/water_sanitation_health/publications/wash-health-care-facilities/en/ [Accessed 22 February 2016]
 23. World Health Organization. Hand hygiene technical reference manual. 2010. Available from: http://www.who.int/gpsc/5may/tools/training_education/en/index.html [Accessed 23 February 2016]
 24. Sax H, Allegranzi B, Uçkay I, Larson E, Boyce J, Pittet D. "My five moments for hand hygiene": a user-centred design approach to understand, train, monitor and report hand hygiene. *J Hosp Infect* 2007; 67:9-21.
 25. Meengs MR et al. Handwashing frequency in an emergency department. *J Emerg Nursing* 1994; 20:183-188.
 26. Zimakoff J, Stormark M, Olesen Larsen S. Use of gloves and handwashing behaviour among health care workers in intensive care units. A multicentre investigation in four hospitals in Denmark and Norway. *J Hosp Infect* 1993; 24:63-67.
 27. Picheansanthian W, Chotibang J. Glove utilization in the prevention of cross transmission: a systematic review. *JBI Database System Rev Implement Rep* 2015; 13(4):188-230. doi: 10.11124/jbisrir-2015-1817.
 28. Rock C, Harris AD, Reich NG, Johnson JK, Thom KA. Is handhygiene before putting on nonsterile gloves in the intensive care unit a waste of health care worker time? a randomized controlled trial. *Am J Infect Control* 2013;41:994-996.
 29. Rossoff LJ, Borenstein M, Isenberg HD. Is hand washing really needed in an intensive care unit? *Crit Care Med* 1995;23:1211-1216.
 30. Loveday HP, Lyman S, Singleton J, Wilson J. Clinical glove use: Health care workers' actions and perceptions. *J Hosp Infect* 2014; 86 (2): 110-6.
 31. WHO: Systematic literature review of automated/electronic systems for hand hygiene monitoring. <http://www.who.int/gpsc/5may/automated-hand-hygiene-monitoring.pdf?ua=1> [Accessed 23 February 2016]
 32. Stewardson A., Allegranzi B., Perneger T., Attar H., Pittet D. Testing the WHO Hand Hygiene Self-Assessment Framework for usability and reliability. *J Hosp Infect* 2013; 83(1):30-5. doi: 10.1016/j.jhin.2012.05.017. Epub 2012 Nov 11.
 33. WHO (2012) Hand Hygiene Self-Assessment Framework Global Survey Summary Report WHO:Geneva http://www.who.int/gpsc/5may/summary_report_HHSAF_global_survey_May12.pdf?ua=1 [Accessed 22 February 2016]

IFIC Basic Concepts of Infection Control, 3rd edition, 2016

34. Luangasanatip N, Hongsuwan M, Limmathurotsakul D, et al. Comparative efficacy of interventions to promote hand hygiene in hospital: systematic review and network meta-analysis. *BMJ* 2015; 351:h3728. doi: 10.1136/bmj.h3728.
35. Allegranzi B, Pittet D. Role of hand hygiene in healthcare-associated infection prevention. *J Hosp Infect* 2009; 73:305-15.
36. Schweizer ML, Reisinger HS, Ohl M, et al. Searching for an optimal hand hygiene bundle: a meta-analysis. *Clin Infect Dis* 2014;58:248–259.
37. Talbot TR, Johnson JG, Fergus C, et al. Sustained improvement in hand hygiene adherence: utilizing shared accountability and financial incentives. *Infect Control Hosp Epidemiol* 2013;34: 1129–1136.
38. Rodriguez V, Giuffr  C, Griselda Almada S, et al. A multimodal intervention to improve hand hygiene in ICUs in Buenos Aires, Argentina: a stepped wedge trial *Inter J Quality Health Care* 2015; 27(5), 405–411.
39. De Bono S, Heling G, Borg MA. Organizational culture and its implications for infection prevention and control in healthcare institutions *J Hosp Infect* 2014; 86(1):1-6.
40. Huis A, van Achterberg T, de Bruin M, et al. A systematic review of hand hygiene improvement strategies: a behavioural approach *Implement Sci* 2012; 14;7:92.
41. Storr J, Wigglesworth N, Kilpatrick C (2013) Integrating human factors with infection prevention and control The Health Foundation: London.
42. Shekelle P., Pronovost P., Wachter R, et al. The Top Patient Safety Strategies That Can Be Encouraged for Adoption Now. *Ann Int Med* 2013; 158 (5) Part 2: 365-369.

Key Web Sites

1. Centers for Disease Control and Prevention: Hand Hygiene in Healthcare Settings. <http://www.cdc.gov/HandHygiene/index.html> [Accessed 23 February 2016]
2. Germs. Wash your hands of them: <http://www.washyourhandsofthem.com/home.aspx> [Accessed 23 February 2016]
3. Institute for Healthcare Improvement (IHI): Improving Hand Hygiene. <http://www.ihl.org/resources/Pages/Tools/HowtoGuidelImprovingHandHygiene.aspx> [Accessed 23 February 2016]
4. 19th WHO Model List of Essential Medicines (April 2015) http://www.who.int/medicines/publications/essentialmedicines/EML2015_8-May-15.pdf [Accessed 23 February 2016]
5. World Health Organization. WHO Starter Toolkit. 2010. <http://www.who.int/gpsc/5may/tools/en/index.html> [Accessed 23 February 2016]

While the advice and information in this chapter is believed to be true and accurate, neither the authors nor the International Federation of Infection Control can accept any legal responsibility or liability for any loss or damage arising from actions or decisions based on this chapter.

Published by the International Federation Of Infection Control
47 Wentworth Green
Portadown, BT62 3WG, N Ireland, UK
www.theifc.org

©International Federation of Infection Control, 2016. All rights reserved.

Table 10.1. World Health Organization Consensus Recommendations Hand Hygiene in Health Care, 2009¹

Recommendation by topic and grade according to the HICPAC ranking system*
<p>1. Indications for hand hygiene</p> <p>A. Wash hands with soap and water when visibly dirty or visibly soiled with blood or other body fluids (IB) or after using the toilet (II).</p> <p>B. If exposure to potential spore-forming pathogens is strongly suspected or proven, including outbreaks of <i>C. difficile</i>, hand washing with soap and water is the preferred means (IB).</p> <p>C. Use an alcohol-based handrub as the preferred means for routine hand antisepsis in all other clinical situations described in items D(a) to D(f) listed below if hands are not visibly soiled (IA). If alcohol-based handrub is not obtainable, wash hands with soap and water (IB).</p> <p>D. Perform hand hygiene:</p> <ul style="list-style-type: none"> a) before and after touching the patient (IB); b) before handling an invasive device for patient care, regardless of whether or not gloves are used (IB); c) after contact with body fluids or excretions, mucous membranes, non-intact skin, or wound dressings (IA); d) if moving from a contaminated body site to another body site during care of the same patient (IB); e) after contact with inanimate surfaces and objects (including medical equipment) in the immediate vicinity of the patient (IB); f) after removing sterile (II) or non-sterile gloves (IB). <p>E. Before handling medication or preparing food perform hand hygiene using an alcohol-based handrub or wash hands with either plain or antimicrobial soap or water (IB).</p> <p>F. Soap and alcohol-based handrub should not be used concomitantly (II).</p>
<p>2. Hand hygiene technique</p> <p>A. Apply a palmful of alcohol-based handrub and cover all surfaces of the hands. Rub hands until dry (IB).</p> <p>B. When washing hands with soap and water, wet hands with water and apply the amount of product necessary to cover all surfaces. Rinse hands with water and dry thoroughly with a single-use towel. Use clean, running water whenever possible. Avoid using hot water, as repeated exposure to hot water may increase the risk of dermatitis (IB). Use a towel to turn off tap/faucet (IB). Dry hands thoroughly using a method that does not recontaminate hands. Make sure towels are not used multiple times or by multiple people (IB).</p> <p>C. Liquid, bar, leaf or powdered forms of soap are acceptable. When bar soap is used, small bars of soap in racks that facilitate drainage should be used to allow the bars to dry (II).</p>
<p>3. Recommendations for surgical hand preparation</p> <p>A. Remove rings, wrist-watch, and bracelets before beginning surgical hand preparation (II). Artificial nails are prohibited (IB).</p> <p>B. Sinks should be designed to reduce the risk of splashes (II).</p> <p>C. If hands are visibly soiled, wash hands with plain soap before surgical hand preparation (II). Remove debris from underneath fingernails using a nail cleaner, preferably under running water (II).</p> <p>D. Brushes are not recommended for surgical hand preparation (IB).</p> <p>E. Surgical hand antisepsis should be performed using either a suitable antimicrobial soap or suitable alcohol-based handrub, preferably with a product ensuring sustained activity, before donning sterile gloves (IB).</p> <p>F. If quality of water is not assured (as described in Table I.11.3) in the operating theatre, surgical hand antisepsis using an alcohol-based handrub is recommended before donning sterile gloves when performing surgical procedures (II).</p> <p>G. When performing surgical hand antisepsis using an antimicrobial soap, scrub hands and forearms for the length of time recommended by the manufacturer, typically 2–5 minutes. Long scrub times (e.g. 10 minutes) are not necessary (IB).</p> <p>H. When using an alcohol-based surgical handrub product with sustained activity, follow the manufacturer's instructions for application times. Apply the product to dry hands only (IB). Do not combine surgical hand scrub and surgical handrub with alcohol-based products sequentially (II).</p> <p>I. When using an alcohol-based handrub, use sufficient product to keep hands and forearms wet with the handrub throughout the surgical hand preparation procedure (IB).</p> <p>J. After application of the alcohol-based handrub as recommended, allow hands and forearms to dry thoroughly before donning sterile gloves (IB).</p>

Table 10.1. World Health Organization Consensus Recommendations Hand Hygiene in Health Care, 2009¹

Recommendation by topic and grade according to the HICPAC ranking system*	
4. Selection and handling of hand hygiene agents	<p>A. Provide HCWs with efficacious hand hygiene products that have low irritancy potential (IB).</p> <p>B. To maximize acceptance of hand hygiene products by HCWs, solicit their input regarding the skin tolerance, feel, and fragrance of any products under consideration (IB). Comparative evaluations may greatly help in this process.</p> <p>C. When selecting hand hygiene products:</p> <ol style="list-style-type: none"> a. determine any known interaction between products used to clean hands, skin care products and the types of glove used in the institution (II); b. solicit information from manufacturers about the risk of product contamination (IB); c. ensure that dispensers are accessible at the point of care (see Part I.1 of the <i>Guidelines</i> for the definition) (IB); d. ensure that dispensers function adequately and reliably and deliver an appropriate volume of the product (II); e. ensure that the dispenser system for alcohol-based handrubs is approved for flammable materials (IC); f. solicit and evaluate information from manufacturers regarding any effect that hand lotions, creams or alcohol-based handrubs may have on the effects of antimicrobial soaps being used in the institution (IB); g. cost comparisons should only be made for products that meet requirements for efficacy, skin tolerance, and acceptability (II). <p>D. Do not add soap (IA) or alcohol-based formulations (II) to a partially empty soap dispenser. If soap dispensers are reused, follow recommended procedures for cleansing.</p>
5. Skin care	<p>A. Include information regarding hand-care practices designed to reduce the risk of irritant contact dermatitis and other skin damage in education programmes for HCWs (IB).</p> <p>B. Provide alternative hand hygiene products for HCWs with confirmed allergies or adverse reactions to standard products used in the healthcare setting (II).</p> <p>C. Provide HCWs with hand lotions or creams to minimize the occurrence of irritant contact dermatitis associated with hand antisepsis or handwashing (IA).</p> <p>D. When alcohol-based handrub is available in the healthcare facility for hygienic hand antisepsis, the use of antimicrobial soap is not recommended (II).</p> <p>E. Soap and alcohol-based handrub should not be used concomitantly (II).</p>
6. Use of gloves	<p>A. The use of gloves does not replace the need for hand hygiene by either handrubbing or handwashing (IB).</p> <p>B. Wear gloves when it can be reasonably anticipated that contact with blood or other potentially infectious materials, mucous membranes or non-intact skin will occur (IC).</p> <p>C. Remove gloves after caring for a patient. Do not wear the same pair of gloves for the care of more than one patient (IB).</p> <p>D. When wearing gloves, change or remove gloves during patient care if moving from a contaminated body site to either another body site (including non-intact skin, mucous membrane or medical device) within the same patient or the environment (II).</p> <p>E. The reuse of gloves is not recommended (IB). In the case of glove reuse, implement the safest re-processing method (II).</p>
7. Other aspects of hand hygiene	<p>A. Do not wear artificial fingernails or extenders when having direct contact with patients (IA).</p> <p>B. Keep natural nails short (tips less than 0.5 cm long or approximately ¼ inch) (II).</p>

Table 10.1. World Health Organization Consensus Recommendations Hand Hygiene in Health Care, 2009¹

Recommendation by topic and grade according to the HICPAC ranking system*	
8.	Educational and motivational programmes for healthcare workers
A.	In hand hygiene promotion programmes for HCWs, focus specifically on factors currently found to have a significant influence on behaviour and not solely on the type of hand hygiene products. The strategy should be multifaceted and multimodal and include education and senior executive support for implementation. (IA).
B.	Educate HCWs about the type of patient-care activities that can result in hand contamination and about the advantages and disadvantages of various methods used to clean their hands (II).
C.	Monitor HCWs' adherence to recommended hand hygiene practices and provide them with performance feedback (IA).
D.	Encourage partnerships between patients, their families and HCWs to promote hand hygiene in health care settings (II).
9.	Governmental and institutional responsibilities
9.1	For healthcare administrators
A.	It is essential that administrators ensure that conditions are conducive to the promotion of a multifaceted, multimodal hand hygiene strategy and an approach that promotes a patient safety culture by implementation of points B–I below.
B.	Provide HCWs with access to a safe, continuous water supply at all outlets and access to the necessary facilities to perform handwashing (IB).
C.	Provide HCWs with a readily accessible alcohol-based handrub at the point of patient care (IA).
D.	Make improved hand hygiene adherence (compliance) an institutional priority and provide appropriate leadership, administrative support, financial resources and support for hand hygiene and other infection prevention and control activities (IB).
E.	Ensure that HCWs have dedicated time for infection control training, including sessions on hand hygiene (II).
F.	Implement a multidisciplinary, multifaceted and multimodal programme designed to improve adherence of HCWs to recommended hand hygiene practices (IB).
G.	With regard to hand hygiene, ensure that the water supply is physically separated from drainage and sewerage within the healthcare setting and provide routine system monitoring and management (IB).
H.	Provide strong leadership and support for hand hygiene and other infection prevention and control activities (II).
I.	Alcohol-based handrub production and storage must adhere to the national safety guidelines and local legal requirements (II).
9.2	For national governments
A.	Make improved hand hygiene adherence a national priority and consider provision of a funded, coordinated implementation programme while ensuring monitoring and long-term sustainability (II).
B.	Support strengthening of infection control capacities within healthcare settings (II).
C.	Promote hand hygiene at the community level to strengthen both self-protection and the protection of others (II).
D.	Encourage healthcare settings to use hand hygiene as a quality indicator (II).

***Ranking system used to grade the recommendations** (Healthcare Infection Control Practices Advisory Committee [HICPAC] of the US Centers for Disease Control and Prevention [CDC]): IA= Strongly recommended for implementation and strongly supported by well-designed experimental, clinical or epidemiological studies. IB=Strongly recommended for implementation and supported by some experimental, clinical or epidemiological studies and a strong theoretical rationale. IC= Required for implementation as mandated by federal and/or state regulation or standard. II=Suggested for implementation and supported by suggestive clinical or epidemiological studies or a theoretical rationale or the consensus of a panel of experts